

Immunotherapy Shows Promise for People With Advanced Anal Cancer

Keytruda may prolong survival compared with standard therapy.

February 6, 2020 By [Liz Highleyman](#)

The checkpoint inhibitor Keytruda (pembrolizumab) demonstrated encouraging overall survival and manageable side effects for people with inoperable or metastatic anal cancer, according to a report at the recent American Society of Clinical Oncology Genitourinary Cancers Symposium in San Francisco.

Although just 12 out of 112 participants in this Phase II clinical trial experienced complete or partial tumor shrinkage, those who did had durable responses lasting nearly three years.

Anal cancer, usually caused by human papillomavirus (HPV), is uncommon, but [new cases and related deaths have risen rapidly](#) in the United States in recent years. Gay and bisexual men—especially those living with HIV—have the highest rate of anal cancer. Standard therapy involves surgery, radiation and chemotherapy, but outcomes are typically poor.

Aurélien Marabelle MD, PhD, of Gustave Roussy cancer center near Paris, presented results from a cohort of patients with anal cancer in KEYNOTE-158, a study evaluating the safety and efficacy of Keytruda in people with various types of solid tumors.

Keytruda is a monoclonal antibody that helps the immune system fight cancer. It blocks PD-1, a checkpoint receptor on T cells that helps regulate immune function. Some tumors can hijack PD-1 to turn off immune responses. Drugs that block the interaction between PD-1 and its binding partner, known as PD-L1, can release the brakes and restore T-cell activity. People with higher levels of PD-L1 in their tumors tend to do better on Keytruda, although this is not a reliable predictor of individual response.

The anal cancer cohort included 112 participants. Just over 80% were women, and the median age was 61. More than 90% had metastatic cancer that had spread elsewhere in the body. About two thirds were classified as having PD-L1 positive tumors. They had previously received at least one prior medication regimen, with about 40% having tried three or more regimens; over 90% had received radiation therapy.

Participants in this nonrandomized, single-arm study received Keytruda administered by IV

infusion every three weeks for up to 35 cycles or until they experienced disease progression or intolerable side effects.

Over a median follow-up period of 12 months, the overall response rate was 10.7%. Six patients (5.4%) achieved complete remission, and six (5.4%) had partial tumor shrinkage. An additional 15.2% had stable disease, but two thirds experienced cancer progression.

Results were a bit better among the 75 participants with PD-L1 positive tumors. This subgroup had an overall response rate of 14.7%, a complete response rate of 8.0%, a partial response rate of 6.7% and a stable disease rate of 14.7%. But those with PD-L1 negative cancer did worse, with only one person (3.3%) seeing a partial response.

Although the proportion of patients who responded was low, those who did so saw a durable benefit, with 90.0% still responding at the one-year and two-year marks.

The median time to disease progression was two months. At the six-month mark, 18.9% were still alive without disease progression, falling to 15.0% at one year.

The median overall survival duration was 11.9 months, with 49.1% still alive after one year on treatment and 25.0% still alive after two years.

Keytruda was generally safe, although side effects were common, experienced by 60.7% of participants. The most frequent adverse events were fatigue, diarrhea, low thyroid function and nausea. Twenty people (17.9%) experienced severe (Grade 3 or 4) adverse events, and five people (4.5%) stopped treatment due to side effects. Three quarters of participants died during the study, underlining the aggressive nature of this cancer.

Unleashing T cells with checkpoint inhibitors can lead to a strong immune response that also harms healthy organs. In this study, 23.2% of participants experienced immune-mediated adverse events or infusion reactions, including five (4.5%) with severe side effects and three (2.7%) who stopped therapy for this reason. The most common immune-mediated events were underactive thyroid function (hypothyroidism), overactive thyroid function (hyperthyroidism) and lung inflammation.

Based on these findings, the researchers concluded that Keytruda demonstrated antitumor activity, durable response and encouraging overall survival with manageable safety in heavily pretreated patients with advanced anal cancer.

[Click here](#) to read the study abstract.

[Click here](#) to learn more about anal cancer.