



# SARS-CoV-2 Antigen Levels Linked to Patient Outcomes

Higher levels of viral antigens in the blood correlated with more severe disease.

September 9, 2022 By National Institutes of Health

---

The amount of SARS-CoV-2 antigen measured in the blood of patients hospitalized with COVID-19 is associated with illness severity and other clinical outcomes, according to a new study [published in the \*Annals of Internal Medicine\*](#).

Following the [ACTIV-3 trial](#) of COVID-19 therapeutics in people hospitalized with COVID-19, researchers from the National Institute of Allergy and Infectious Diseases (NIAID), part of the National Institutes of Health, and their collaborators analyzed levels of SARS-CoV-2 antigen in blood samples taken from study participants and assessed the association of those levels with disease progression.

The authors suggest that SARS-CoV-2 antigen levels hold promise as a biomarker, or a measurable substance, to predict which patients hospitalized with COVID-19 have a higher risk of worse outcomes.

The ACTIV-3 trial enrolled people hospitalized with COVID-19 between August 2020 and November 2021. Participants contributed a baseline blood sample and were then randomized to receive either an experimental COVID-19 therapeutic or a placebo. All participants received the antiviral remdesivir unless contraindicated. In this follow-up analysis, the researchers examined 2540 participant baseline blood samples for SARS-CoV-2 antigen levels.

The researchers assessed the relationship between each participant's SARS-CoV-2 blood antigen levels and their time to discharge from the hospital, as well as their pulmonary symptoms at Day 5 of the trial—whether they had stayed the same, worsened, or improved since enrollment. With all this information in hand, the researchers conducted statistical analyses to determine if plasma antigen levels were associated with the participants' pulmonary function when they gave the blood sample—and whether they could predict how the participants would fare over time. In addition, the researchers examined the relationship between a number of participant and viral characteristics and antigen levels.

The analysis revealed a strong correlation between higher SARS-CoV-2 antigen levels ( $\geq 1000$  nanograms per liter) and worse pulmonary function at the time of enrollment. Importantly,

participants with higher SARS-CoV-2 antigen levels in the blood at enrollment generally had decreased pulmonary function at Day 5, regardless of the severity of their illness at the time of study entry, and took longer to be released from the hospital. High blood antigen levels also correlated with some known risk factors for worse illness, such as being male.

Three additional participant characteristics were found to correlate with lower antigen levels: the presence of SARS-CoV-2 antibodies, exposure to remdesivir prior to enrollment and longer time in hospital prior to enrollment. Finally, participants infected with the delta variant had higher antigen levels than those infected with prior circulating strains.

The researchers concluded that antigen levels in the blood are likely a marker of ongoing viral replication and could be useful for predicting a patient's disease progression and likely outcomes once they are admitted to the hospital. The authors note that these results suggest that a precision medicine approach could be helpful in future clinical trials of antiviral therapy. For instance, antigen levels could help determine those patients most likely to benefit from therapies directed at fighting the virus.

This news release was originally published by the National Institutes of Health on August 29, 2022.

---

© 2026 Smart + Strong All Rights Reserved.

<http://beta.docker.poz.com/article/sarscov2-antigen-levels-linked-patient-outcomes>