



Studying HIV Among Women in the South

Two projects in Alabama explore why they're more vulnerable.

June 26, 2023 By Laura Schmidt

Why are HIV rates among women in the South higher than in other regions—and what can be done to address this health disparity? The National Institutes of Health (NIH) awarded researchers at the University of Alabama at Birmingham (UAB) a combined \$19.5 million in federal grants to investigate these very questions, according to a UAB press release.

Although 38% of the U.S. population lives in the South, the region accounts for 53% of new HIV cases and 47% of all deaths among people with HIV, according to the Centers for Disease Control and Prevention. Specifically, it's estimated that 18,500 of the 34,800 new HIV diagnoses in 2019 were in the South. What's more, about 55% of women diagnosed with HIV in the United States live in the South, according to AIDSVu.

The NIH grants will fund two studies focusing on the ways behavioral, geographic and demographic factors relate to HIV rates among women in the South, according to the press release.

The bulk of the funding—\$15 million—will go to the AWARE project, which is led by researchers at UAB, the Columbia School of Nursing in New York City and Lurie Children's Hospital in Chicago.

The AWARE Project will enroll a national digital cohort of 1,800 HIV-negative women to study how various factors impact the risk of acquiring HIV as well as other sexually transmitted infections (STIs).

The second study, the Camellia Project, will receive \$4.5 million and be co-led by Latesha Elope, MD, an associate professor in the UAB Division of Infectious Diseases in the Marnix E. Heersink School of Medicine, and Lynn Matthews, MD, an associate professor in the UAB Division of Infectious Diseases.

In addition to HIV and STIs, the Camellia Project will also focus on access and use of pre-exposure prophylaxis (PrEP) among vulnerable cis- and transgender women in Alabama. PrEP refers to the daily pills or long-acting injectables that prevent a person from contracting HIV.

“The rate of HIV infection is 10 times higher for Black women compared to white women; however, PrEP utilization remains low,” Elope says.

The study will identify the barriers that prevent people of color, specifically women in the Deep South, from accessing PrEP.

© 2026 Smart + Strong All Rights Reserved.

<http://beta.docker.poz.com/article/studying-hiv-among-women-south>