



Staying On Track

June 25, 2014 By Jennifer Morton

Adherence—taking your medication as scheduled—is essential to treating HIV. Missing doses can cause drug resistance and limit your future treatment options. If you’re looking to improve your adherence, or if you’re preparing to start a new regimen, it’s important to discuss any fears or challenges you might face with your health care provider. We asked you about your HIV adherence and what you do to make sure you take your meds on time.

In the past month, how many times did you miss a dose of your HIV meds?

- 64% - Never
- 18% - 1
- 9% - 2
- 9% - 3 or more

Do you use a reminder to take your HIV meds?

- 45% - Yes
- 55% - No

Do you carry or store your meds in a special case?

- 63% - Yes
- 37% - No

Have you ever...

...missed several doses in a row?

- 17% - Yes
- 83% - No

...taken a double dose by mistake?

- 18% - Yes
- 82% - No

...run out of your HIV meds?

- 20% - Yes
- 80% - No

...switched your meds to improve adherence?

24% - Yes

76% - No

Adherence Tip: Keep your meds next to something you use on a daily basis like your coffeepot, your alarm clock or your toothbrush to help you remember to take them each day.

Source: October/November 2013

© 2026 Smart + Strong All Rights Reserved.

<http://beta.docker.poz.com/article/survey-says-july-august-2014-25814-7819>