

Transmitted HIV Drug Resistance Is Rising, Yet Isn't a Major Threat to Treatment

Most of such drug resistance seen is to older drugs.

June 24, 2018 By [Benjamin Ryan](#)

The rate of people acquiring HIV that is resistant to antiretrovirals (ARVs) has risen in recent years, [aidsmap](#) reports. However, because for the most part, such strains of the virus have mutations that confer resistance to older HIV medications, their spread is not likely a significant impediment to finding an effective drug regimen.

Publishing their findings in *Clinical Infectious Diseases*, researchers at Kaiser Permanente Northern California studied 2003 to 2015 data on 4,353 people with HIV who were patients of the integrated health care system, received genotypic HIV drug resistance testing and initially had not taken ARVs.

Nearly 14 percent of the cohort had acquired drug-resistant HIV, most commonly virus that was resistant to non-nucleoside reverse transcriptase inhibitors, or NNRTIs (7.2 percent of the cohort), followed by virus resistant to nucleoside/nucleotide reverse transcriptase inhibitors, or NRTIs (5.8 percent), protease inhibitors (3.2 percent) and multiple drug classes (1.9 percent).

There were no reports of virus resistant to the newest class of ARVs, integrase inhibitors, which are well established as posing a low risk of drug resistance development.

Each year, a median 305 people in the cohort received resistance testing. During the study period, the proportion of people who had acquired HIV that was resistant to any type of ARV increased by 5 percent per year, including 11 percent per year for NNRTIs.

The ARVs to which such viral strains were most commonly resistant included Sustiva (efavirenz) and Viramune (nevirapine). Nearly 8 percent of the entire cohort had some level of resistance to Sustiva. Other drugs to which people acquired drug resistance included Edurant (rilpivirine) (5 percent of the cohort), Ziagen (abacavir) (2.8 percent), Viread (tenofovir disoproxil fumarate, or TDF) (2.2 percent), Reyataz (atazanavir) and Norvir (ritonavir)-boosted Prezista (darunavir) (0.3 percent).

Based on the fact that there was scant evidence of HIV that transmitted with resistance to newer drugs, the researchers theorized that the transmitted drug resistance seen in the study was largely acquired by cohort members from individuals who themselves had acquired HIV that was already drug resistant.

To read the aidsmap article, [click here](#).

To read the study abstract, [click here](#).

© 2026 Smart + Strong All Rights Reserved.

<http://beta.docker.poz.com/article/transmitted-hiv-drug-resistance-rising-yet-major-threat-treatment>